

CORRECTION OF “ABERRANT PLUMAGE IN A BLACK-AND-WHITE TANAGER (*CONOTHTRAUPIS SPECULIGERA*)”

Daniel J. Lebbin

Department of Ecology & Evolutionary Biology, Cornell University, E148 Corson
Hall, Ithaca, NY 14853

djl42@cornell.edu

In Lebbin (2005), I erroneously described a semiconcealed white crown patch in a male Black-and-white Tanager (*Conothraupis speculigera*) captured in Peru as an aberrant plumage character for this species. Although not described nor illustrated by Isler & Isler (1999) and Ridgely & Greenfield (2001), the semiconcealed white crown patch was described by Witt (2005). All adult male *C. speculigera* specimens at Louisiana State University Museum of Natural Science (n = 4; LSUMZ 31456, 113916, 113929, 172388) show this semiconcealed white crown patch. This white patch is absent on all non-adult male *C. speculigera* skins (females n = 14; “female?” n = 1, immature males n = 16), including 5 immature males showing mostly single black feathers on the lores, malar, or sides of breast, but not on

the crown or nape (LSUMZ 28638, 113919, 113926, 113932, 113941). Upon consulting with specimen material, it is clear that this plumage feature is not aberrant and had been previously described. I must have overlooked this plumage feature when handling the first male (which I described as not having the concealed white patch in Lebbin 2005) at Playa Bonita.

Acknowledgements

D. F. Lane provided useful comments on this correction. D. Calderón-F. and J. P. O’Neill provided useful feedback after the original manuscript was published. J. V. Remsen Jr., S. W. Cardiff, and D. F. Lane provided useful data from the LSUMZ collection.

References

Isler, M. L. & P. R. Isler. 1999. *The Tanagers: Natural History, Distribution, and Identification*. Smithsonian Institution Press, Washington, DC.

- Lebbin, D. J. 2005. Aberrant plumage in a black-and-white tanager (*Conothraupis speculigera*). Boletín SAO. 15(2): 100-104.
- Ridgely, R. S. & P. J. Greenfield. 2001. The birds of Ecuador: field guide. Cornell University Press, Ithaca, New York.
- Witt, C. C. 2005. Syntopic breeding suggests mimicry of the Black-and-white Seedeater (*Sporophila luctuosa*) by the Black-and-White Tanager (*Conothraupis speculigera*). Ornitologia Neotropical 16(3): 387-396.